

Post-operative Instructions Following Minor Ocular Surgery

Patient Name: _____

Date: _____

You have just received an injection of medication into the _____ eye.

It is important to review the following items:

- Use _____ (Tan Cap) one drop 4 times a day for 7 days in the injected eye.
- Shake the eye drops before each use and close your eyes for 2 minutes after each drop.
- Do not rub or apply pressure to your eye. ● Avoid getting water into the eye.
- **Redness** may be present on the side of your eye where the injection was placed.
This should go away over the next week. However, if you notice new or increased redness beginning in the eye, please call immediately.
- **Mild Pain** or **Irritation** may occur this evening and tomorrow.
Please use Tylenol for this pain. You may use one or two drops of artificial tears as needed to lubricate the eye. If the pain continues, or you develop new pain during the week, please call immediately.
- **Floaters** may be noticed in your vision over the next few days.
This is the medication and it is normal to see a few floaters following this type of injection. The floaters will settle as the medication absorbs. If you develop a sudden shower of many floaters or new flashes of light, please call immediately.
- **Blurry Vision** may be present this evening and tomorrow.
As the medication begins to absorb your baseline vision should recover. If you notice a loss of vision over the next week, please call immediately.
- **In Case of Emergency** call our office immediately.
You may also call anytime with other questions or concerns. Our staff will contact you in 1 week to ensure that you are doing well. In the event that you do not hear from us within 1 week, please call to report your status.

It is a pleasure taking care of you.

Jeremiah Brown, Jr., MS, M.D.